

## Higher $(U, M)$ - Derivations in Completely Semi Prime $\Gamma$ -Rings

M.M. Rahman<sup>1</sup> and S.K. Paul<sup>2,\*</sup>

<sup>1</sup>Department of Mathematics, Jagannath University, Dhaka, Bangladesh.

<sup>2,\*</sup>Department of Basic Sciences and Humanities, University of Asia Pacific, Dhaka, Bangladesh.

\*e-mail: sanjitkpaul@gmail.com

*The aim of the present paper is to prove  $d_n(uam) = \sum_{i+j=n} d_i(u)ad_j(m), \forall u \in U, m \in M, \alpha \in \Gamma, n \in \mathbb{N}$ , where  $M$  is a 2-torsion free completely semiprime  $\Gamma$ -ring satisfying the condition  $aab\beta c = a\beta bac, \forall a, b, c \in M$  and  $\alpha, \beta \in \Gamma, U$  is an admissible Lie ideal of  $M$  and  $D = (d_i)_{i \in \mathbb{N}}$  is a higher  $(U, M)$ -derivation of  $M$ .*

**Keywords:** Admissible Lie ideal, higher  $(U, M)$ -derivation, completely semiprime  $\Gamma$ -ring.

**2010 AMS Subject Classification:** Primary 13N15; Secondary 16W10, 17C50.

### 1. INTRODUCTION

If there is a mapping  $M \times \Gamma \times M \rightarrow M$  ( $M$  and  $\Gamma$  are additive abelian groups) such that

$$(i) \quad (x + y)az = xaz + yaz, \quad x(\alpha + \beta)y = x\alpha y + x\beta y, \quad x\alpha(y + z) = x\alpha y + x\alpha z$$

$$(ii) \quad (x\alpha y)\beta z = x\alpha(y\beta z), \quad \forall x, y, z \in M; \quad \alpha, \beta \in \Gamma,$$

then  $M$  is called a  $\Gamma$ -ring. This concept is more general than that of a ring. Barnes [1] generalized the notion of this type of ring which was introduced by Nobusawa [2]. A  $\Gamma$ -ring  $M$  is a *prime  $\Gamma$ -ring* if  $\forall a, b \in M, a\Gamma M\Gamma b = 0$  implies  $a = 0$  or  $b = 0$ ,  $M$  is called *semiprime* if  $a\Gamma M\Gamma a = 0$  (with  $a \in M$ ) implies  $a = 0$  and it will be *completely semiprime* if  $a\Gamma a = 0$  (with  $a \in M$ ) implies  $a = 0$ . A  $\Gamma$ -ring  $M$  is *2-torsion free* if  $2a = 0$  implies  $a = 0, \forall a \in M$ .

For any  $x, y \in M$  and  $\alpha \in \Gamma$ , we induce a new product, the *Lie product* by  $[x, y]_\alpha = x\alpha y - y\alpha x$ . An additive subgroup  $U \subset M$  is said to be a *Lie ideal* of  $M$  if whenever  $u \in U, m \in M$  and  $\alpha \in \Gamma$ , then  $[u, m]_\alpha \in U$ . In the main results of this article we assume that the Lie ideal  $U$  verifies  $u\alpha u \in U, \forall u \in U$ . A Lie ideal of this type is called a *square closed Lie ideal*. Furthermore, if the Lie ideal  $U$  is square closed and  $U$  is not contained in  $Z(M)$ , where  $Z(M)$  denotes the center of  $M$ , then  $U$  is called an *admissible Lie ideal* of  $M$ .

Throughout the article, we use the condition  $aab\beta c = a\beta bac, \forall a, b, c \in M$  and  $\alpha, \beta \in \Gamma$  and this is represented by (\*). We make the basic commutator identities:

$$[x\alpha y, z]_\beta = [x, z]_\beta \alpha y + x[\alpha, \beta]_z y + x\alpha[y, z]_\beta, \quad [x, y\alpha z]_\beta = [x, y]_\beta \alpha z - y[\alpha, \beta]_x z + y\alpha[x, z]_\beta, \\ \forall a, b, c \in M, \forall \alpha, \beta \in \Gamma. \text{ According to the condition (*), the above two identities reduces to}$$

$$[x\alpha y, z]_\beta = [x, z]_\beta \alpha y + x\alpha [y, z]_\beta, [x, y\alpha z]_\beta = [x, y]_\beta \alpha z + y\alpha [x, z]_\beta, \forall a, b, c \in M, \forall \alpha, \beta \in \Gamma.$$

Herstein [3] proved a well known result in prime rings which states that 'every Jordan derivation on a 2-torsion free prime ring is a derivation'. Afterwards many mathematicians studied extensively the derivations in prime rings.  $(U, R)$ -derivations in rings have been introduced by Faraj, Haetinger and Majeed [4], as a generalization of Jordan derivations on a Lie ideal of a ring. The notion of  $(U, R)$ -derivation extends the concept given in [5]. Faraj, Haetinger and Majeed [4] proved that if  $R$  is a prime ring,  $\text{char}(R) \neq 2$ ,  $U$  is a square closed Lie ideal of  $R$  and  $d$  is a  $(U, R)$ -derivation of  $R$ , then  $d(ur) = d(u)r + ud(r), \forall u \in U, r \in R$ .

The notions of derivation and Jordan derivation in  $\Gamma$ -rings have been introduced by Sapanci and Nakajima [6]. Afterwards, in the light of some significant results due to Jordan left derivation of a classical ring obtained by Jun and Kim [7], some extensive results of left derivation and Jordan left derivation of a  $\Gamma$ -ring were determined by Ceven [8]. Halder and Paul [9] extended the results of Lie ideals [8].

In this article, we prove that  $d_n(uam) = \sum_{i+j=n} d_i(u)\alpha d_j(m), \forall u \in U, m \in M, \alpha \in \Gamma, n \in \mathbb{N}$ , where  $M$  is a 2-torsion free completely semiprime  $\Gamma$ -ring satisfying the condition  $aab\beta c = a\beta b\alpha c, \forall a, b, c \in M$  and  $\alpha, \beta \in \Gamma$ ,  $U$  is an admissible Lie ideal of  $M$  and  $D = (d_i)_{i \in \mathbb{N}}$  is a higher  $(U, M)$ -derivation of  $M$ .

## 2. $(U, M)$ -DERIVATION

In this section we recall some definitions and lemmas (established earlier) that are important for representing our main objective in the later section.

Rahman and Paul [10] introduced the concept of  $(U, M)$ -derivation of  $\Gamma$ -ring in the following way:

**Definition 1:** Let  $M$  be a  $\Gamma$ -ring and  $U$  be a Lie ideal of  $M$ . An additive mapping  $d: M \rightarrow M$  is said to be a  $(U, M)$ -derivation of  $M$  if  $d(u\alpha m + s\alpha u) = d(u)\alpha m + u\alpha d(m) + d(s)\alpha u + s\alpha d(u), \forall u \in U, m, s \in M$  and  $\alpha \in \Gamma$ .

**Lemma 2.1:** ([10, Lemma 1]) Let  $M$  be a 2-torsion free  $\Gamma$ -ring satisfying the condition (\*), and  $U$  be a Lie ideal of  $M$  and  $d$  be a  $(U, M)$ -derivation of  $M$ , then

- (i)  $d(u\alpha m\beta u) = d(u)\alpha m\beta u + u\alpha d(m)\beta u + u\alpha m\beta d(u), \forall u \in U, m \in M$  and  $\alpha, \beta \in \Gamma$ .
- (ii)  $d(u\alpha m\beta v + v\alpha m\beta u) = d(u)\alpha m\beta v + u\alpha d(m)\beta v + u\alpha m\beta d(v) + d(v)\alpha m\beta u + v\alpha d(m)\beta u + v\alpha m\beta d(u), \forall u, v \in U, m \in M$  and  $\alpha, \beta \in \Gamma$ .

**Definition 2:** Let  $d$  be a  $(U, M)$ -derivation of  $M$ . We define  $\phi_\alpha(u, m) = d(u\alpha m) - d(u)\alpha m - u\alpha d(m), \forall u \in U, m \in M$  and  $\alpha \in \Gamma$ .

**Lemma 2.2:** Let  $d$  be a  $(U, M)$ -derivation of  $M$ , then  $\forall u, v \in U, m \in M$  and  $\alpha \in \Gamma$ :

- (i)  $\phi_\alpha(u, m) = -\phi_\alpha(m, u)$

(ii)  $\phi_\alpha(u + v, m) = \phi_\alpha(u, m) + \phi_\alpha(v, m), \forall u, v \in U, m \in M$  and  $\alpha \in \Gamma$ .

**Lemma 2.3:** ([10, Lemma 4]) Let  $M$  be a 2-torsion free prime  $\Gamma$ -ring satisfying the condition (\*) and  $U$  be a Lie ideal of  $M$  such that  $U \dot{U}Z(M)$ . Then there exist elements  $a, b \in U$  such that  $[a, b]_\alpha = a\alpha b - b\alpha a \neq 0$ .

**Lemma 2.4:** ([10, Lemma 5]) Let  $M$  be a 2-torsion free prime  $\Gamma$ -ring satisfying the condition (\*) and  $U$  be an admissible Lie ideal of  $M$ . If  $t\alpha v\beta v + v\alpha v\beta t = 0$  for any  $t \in M, v \in U$  and  $\alpha, \beta \in \Gamma$ , then  $t = 0$ .

**Lemma 2.5:** ([10, Theorem 1]) Let  $M$  be a 2-torsion free prime  $\Gamma$ -ring satisfying the condition (\*),  $U$  be an admissible Lie ideal of  $M$  and  $d$  be a  $(U, M)$ -derivation of  $M$ . Then  $\phi_\alpha(u, v) = 0, \forall u, v \in U$  and  $\alpha \in \Gamma$ .

**Lemma 2.6:** ([10, Lemma 6]) Let  $M$  be a 2-torsion free prime  $\Gamma$ -ring satisfying the condition (\*),  $U$  be an admissible Lie ideal of  $M$  and  $d$  be a  $(U, M)$ -derivation of  $M$ . Then  $\phi_\beta(u\alpha u, m) = 0, \forall u \in U, m \in M$  and  $\alpha, \beta \in \Gamma$ .

**Theorem 2.1:** ([10, Theorem 2]) Let  $M$  be a 2-torsion free prime  $\Gamma$ -ring satisfying the condition (\*),  $U$  be a square closed Lie ideal of  $M$  and  $d$  be a  $(U, M)$ -derivation of  $M$ . Then  $d(u\alpha m) = d(u)\alpha m + u\alpha d(m), \forall u \in U, m \in M, \alpha \in \Gamma$ .

**Definition 3.** Let  $M$  be a  $\Gamma$ -ring and  $U$  be a Lie ideal of  $M$  and  $D = (d_i)_{i \in \mathbb{N}_0}$  be a family of additive mappings of  $M$  into itself such that  $d_0 = id_M$ , where  $id_M$  is an identity mapping on  $M$ . Then  $D$  is said to be a higher  $(U, M)$ -derivation of  $M$  if for each  $n \in \mathbb{N}, d_n(u\alpha m + sau) = \sum_{i+j=n} d_i(u)\alpha d_j(m) + d_i(s)\alpha d_j(u), \forall u \in U, m, s \in M$  and  $\alpha, \beta \in \Gamma$ .

**Lemma 2.7:** Let  $M$  be a 2-torsion free  $\Gamma$ -ring satisfying the condition (\*),  $U$  be a Lie ideal of  $M$  and  $D = (d_i)_{i \in \mathbb{N}}$  be a higher  $(U, M)$ -derivation of  $M$ . Then  $d_n(u\alpha m\beta u) = \sum_{i+j+k=n} d_i(u)\alpha d_j(m)\beta d_k(u), \forall u \in U, m \in M$  and  $\alpha, \beta \in \Gamma$ .

**Proof:** Let  $x = u\alpha((2u)\beta m + m\beta(2u)) + ((2u)\beta m + m\beta(2u))\alpha u$ .

We have, by Definition 3

$$d_n(u\alpha m + sau) = \sum_{i+j=n} d_i(u)\alpha d_j(m) + d_i(s)\alpha d_j(u) \quad (A)$$

$$\text{If } s = m, \text{ then } d_n(u\alpha m + mau) = \sum_{i+j=n} d_i(u)\alpha d_j(m) + d_i(m)\alpha d_j(u) \quad (B)$$

Now replacing  $m$  and  $s$  by  $(2u)\beta m + m\beta(2u)$  and  $(2u)\alpha m + m\alpha(2u)$  respectively in (A) and using (B) and the condition (\*), we get

$$\begin{aligned} & d_n(u\alpha((2u)\beta m + m\beta(2u)) + ((2u)\alpha m + m\alpha(2u))\beta u) \\ &= \sum_{i+j=n} d_i(u)\alpha d_j((2u)\beta m + m\beta(2u)) + d_i((2u)\beta m + m\beta(2u))\alpha d_j(u) \end{aligned}$$

This implies that

$$\begin{aligned}
 d_n(x) &= 2 \sum_{i+j=n} d_i(u)\alpha \sum_{l+t=j} (d_l(u)\beta d_t(m) + d_l(m)\beta d_t(u)) \\
 &\quad + 2 \sum_{p+q=i} \sum_{i+j=n} d_j(u)\alpha (d_p(u)\beta d_q(m) + d_p(m)\beta d_q(u)) \\
 &= 2 \sum_{i+l+t=n} (d_i(u)\alpha d_l(u)\beta d_t(m) + d_i(u)\alpha d_l(m)\beta d_t(u)) \\
 &\quad + 2 \sum_{p+q+j=n} (d_p(u)\beta d_q(m)\alpha d_j(u) + d_p(m)\beta d_q(u)\alpha d_j(u)) \\
 &= 2 \sum_{i+l+t=n} (d_i(u)\alpha d_l(u)\beta d_t(m) + d_i(u)\alpha d_l(m)\beta d_t(u)) + \\
 &\quad 2 \sum_{p+q+j=n} (d_p(u)\alpha d_q(m)\beta d_j(u) + d_p(m)\alpha d_q(u)\beta d_j(u)) \tag{1}
 \end{aligned}$$

On the other hand by the definition of higher  $(U, M)$ -derivation and using the condition (\*)

$$\begin{aligned}
 d_n(x) &= 2d_n((u\alpha u)\beta m + m\beta(u\alpha u)) + 2d_n(u\alpha m\beta u) + 2d_n(u\beta m\alpha u) \\
 &= 2d_n((u\alpha u)\beta m + m\beta(u\alpha u)) + 2d_n(u\alpha m\beta u) + 2d_n(u\alpha m\beta u) \\
 &= 2 \sum_{i+j=n} (d_i(u\alpha u)\beta d_j(m) + d_i(m)\beta d_j(u\alpha u)) + 4d_n(u\alpha m\beta u) \\
 &= 2 \sum_{i+j=n} \sum_{r+s=i} d_r(u)\alpha d_s(u)\beta d_j(m) + 2 \sum_{i+j=n} \sum_{e+k=j} d_i(m)\alpha d_e(u)\beta d_k(u) \\
 &\quad + 4d_n(u\alpha m\beta u) \\
 &= 2 \sum_{r+s+j=n} d_r(u)\alpha d_s(u)\beta d_j(m) + 2 \sum_{i+e+k=n} d_i(m)\beta d_e(u)\alpha d_k(u) + 4d_n(u\alpha m\beta u) \tag{2}
 \end{aligned}$$

Now comparing (1) and (2) we get

$$4d_n(u\alpha m\beta u) = 4 \sum_{i+j+k=n} d_i(u)\alpha d_j(m)\beta d_k(u), \forall u \in U, m \in M \text{ and } \alpha, \beta \in \Gamma.$$

Using 2-torsion freeness of  $M$  we get the desired result.

**Lemma 2.8:** Let  $M$  be a 2-torsion free  $\Gamma$ -ring satisfying the condition (\*),  $U$  be a Lie ideal of  $M$  and  $D = (d_i)_{i \in \mathbb{N}}$  be a higher  $(U, M)$ -derivation of  $M$ . Then  $d_n(u\alpha m\beta v + v\alpha m\beta u) = \sum_{i+j+k=n} d_i(u)\alpha d_j(m)\beta d_k(v) + d_i(v)\alpha d_j(m)\beta d_k(u), \forall u, v \in U, m \in M$  and  $\alpha, \beta \in \Gamma$ .

**Proof:** Linearizing of  $d_n(u\alpha m\beta u) = \sum_{i+j+k=n} d_i(u)\alpha d_j(m)\beta d_k(u)$  with respect to  $u$  gives

$$\begin{aligned}
 d_n((u + v)\alpha m\beta(u + v)) &= \sum_{i+j+k=n} d_i(u + v)\alpha d_j(m)\beta d_k(u + v) \\
 &= \sum_{i+j+k=n} (d_i(u)\alpha d_j(m)\beta d_k(u) + d_i(u)\alpha d_j(m)\beta d_k(v) \\
 &\quad + d_i(v)\alpha d_j(m)\beta d_k(u) + d_i(v)\alpha d_j(m)\beta d_k(v)) \tag{3}
 \end{aligned}$$

On the other hand

$$\begin{aligned}
 d_n((u + v)\alpha m\beta(u + v)) &= d_n(u\alpha m\beta u) + d_n(u\alpha m\beta v + v\alpha m\beta u) + d_n(v\alpha m\beta v) \\
 &= \sum_{i+j+k=n} (d_i(u)\alpha d_j(m)\beta d_k(u) + d_n(u\alpha m\beta v + v\alpha m\beta u) \\
 &\quad + \sum_{i+j+k=n} (d_i(v)\alpha d_j(m)\beta d_k(v)) \tag{4}
 \end{aligned}$$

Now comparing (3) and (4) we get

$$d_n(uam\beta v + v\alpha m\beta u) = \sum_{i+j+k=n} d_i(u)\alpha d_j(m)\beta d_k(v) + d_i(v)\alpha d_j(m)\beta d_k(u), \forall u, v \in U, m \in M \text{ and } \alpha, \beta \in \Gamma.$$

**Definition 4:** Let  $M$  be a 2-torsion free  $\Gamma$ -ring satisfying the condition (\*) and  $U$  be a Lie ideal of  $M$ . For every higher  $(U, M)$ -derivation  $D = (d_i)_{i \in \mathbb{N}}$  of  $M$ , we define

$$\phi_n^\alpha(u, m) = d_n(uam) - \sum_{i+j=n} d_i(u)\alpha d_j(m), \forall u \in U, m \in M, \alpha \in \Gamma \text{ and } n \in \mathbb{N}.$$

**Lemma 2.9:** Let  $M$  be a 2-torsion free  $\Gamma$ -ring satisfying the condition (\*) and  $U$  be a Lie ideal of  $M$ . For every  $u \in U, m \in M, \alpha \in \Gamma$  and  $n \in \mathbb{N}$ , then  $\phi_n^\alpha(u, m) + \phi_n^\alpha(m, u) = 0$ .

**Lemma 2.10:** Let  $M$  be a 2-torsion free  $\Gamma$ -ring satisfying the condition (\*) and let  $D = (d_i)_{i \in \mathbb{N}}$  be a higher  $(U, M)$ -derivation of  $M$ . Let  $n \in \mathbb{N}$  and assume that  $a, b \in U, \alpha, \beta, \gamma \in \Gamma$ . If  $\phi_p^\alpha(a, b) = 0$ , for every  $p < n$ , then  $\phi_n^\alpha(a, b) = 0, \forall a, b \in U, \alpha \in \Gamma$  and  $n \in \mathbb{N}$ .

**Proof:** Let  $G = d_n(aab\beta[a, b]_\alpha\gamma b\alpha a + b\alpha a\beta[a, b]_\alpha\gamma aab)$ .

First we compute  $G = d_n(\alpha a(b\beta[a, b]_\alpha\gamma b)\alpha a) + d_n(b\alpha(a\beta[a, b]_\alpha\gamma a)\alpha b)$ .

Using Lemma 2.7, we have

$$\begin{aligned} G &= \sum_{i+f+l=n} d_i(a)\alpha d_f(b\beta[a, b]_\alpha\gamma b)\alpha d_l(a) + \sum_{i+f+l=n} d_i(b)\alpha d_f(a\beta[a, b]_\alpha\gamma a)\alpha d_l(b) \\ &= \sum_{i+j+k+h+l=n} d_i(a)\alpha d_j(b)\beta d_k([a, b]_\alpha)\gamma d_h(b)\alpha d_l(a) \\ &\quad + \sum_{i+j+k+h+l=n} d_i(b)\alpha d_j(a)\beta d_k([a, b]_\alpha)\gamma d_h(a)\alpha d_l(b) \end{aligned}$$

On the other hand

$$G = d_n((aab)\beta[a, b]_\alpha\gamma(b\alpha a) + (b\alpha a)\beta[a, b]_\alpha\gamma(aab)).$$

Using Lemma 2.8, we obtain

$$G = \sum_{r+s+t=n} d_r(aab)\beta d_s([a, b]_\alpha)\gamma d_t(b\alpha a) + \sum_{r+s+t=n} d_r(b\alpha a)\beta d_s([a, b]_\alpha)\gamma d_t(aab).$$

Comparing both expression of  $G$ , we obtain

$$\begin{aligned} &\sum_{i+j+k+h+l=n} d_i(a)\alpha d_j(b)\beta d_k([a, b]_\alpha)\gamma d_h(b)\alpha d_l(a) + \\ &\sum_{i+j+k+h+l=n} d_i(b)\alpha d_j(a)\beta d_k([a, b]_\alpha)\gamma d_h(a)\alpha d_l(b) = \\ &\sum_{r+s+t=n} d_r(aab)\beta d_s([a, b]_\alpha)\gamma d_t(b\alpha a) + \sum_{r+s+t=n} d_r(b\alpha a)\beta d_s([a, b]_\alpha)\gamma d_t(aab) \end{aligned}$$

This implies

$$\sum_{i+j+k+h+l=n} d_i(a)\alpha d_j(b)\beta d_k([a, b]_\alpha)\gamma d_h(b)\alpha d_l(a) -$$

$$\begin{aligned} \sum_{r+s+t=n} d_r(aab)\beta d_s([a, b]_\alpha)\gamma d_t(baa) = \\ -[\sum_{i+j+k+h+l=n} d_i(b)\alpha d_j(a)\beta d_k([a, b]_\alpha)\gamma d_h(a)\alpha d_l(b) - \\ \sum_{r+s+t=n} d_r(baa)\beta d_s([a, b]_\alpha)\gamma d_t(aab)] \end{aligned} \tag{5}$$

By the inductive assumption we can put  $d_r(x\alpha y)$  for  $\sum_{i+j=r} d_i(x)\alpha d_j(y)$ , when  $r < n$ , for  $x = a, b$  and  $y = b, a$ .

Thus, we can write by Definition 4

$$\begin{aligned} \sum_{i+j+k+h+l=n} (d_i(a)\alpha d_j(b)\beta d_k([a, b]_\alpha)\gamma d_h(b)\alpha d_l(a)) \\ - \sum_{r+s+t=n} (d_r(aab)\beta d_s([a, b]_\alpha)\gamma d_t(baa)) \\ = -(\phi_n^\alpha(a, b)\beta[a, b]_\alpha\gamma baa + aab\beta[a, b]_\alpha\gamma\phi_n^\alpha(b, a)) \end{aligned} \tag{6}$$

Now using (6), (5) becomes

$$\phi_n^\alpha(a, b)\beta[a, b]_\alpha\gamma baa + aab\beta[a, b]_\alpha\gamma\phi_n^\alpha(b, a) + \phi_n^\alpha(b, a)\beta[a, b]_\alpha\gamma aab + baa\beta[a, b]_\alpha\gamma\phi_n^\alpha(a, b) = 0.$$

By Lemma 2.4, we have

$$\begin{aligned} \phi_n^\alpha(a, b)\beta[a, b]_\alpha\gamma baa - aab\beta[a, b]_\alpha\gamma\phi_n^\alpha(a, b) - \phi_n^\alpha(a, b)\beta[a, b]_\alpha\gamma aab \\ + baa\beta[a, b]_\alpha\gamma\phi_n^\alpha(a, b) = 0. \end{aligned}$$

This implies that

$$\phi_n^\alpha(a, b)\beta[a, b]_\alpha\gamma[a, b]_\alpha + [a, b]_\alpha\beta[a, b]_\alpha\gamma\phi_n^\alpha(a, b) = 0.$$

Thus by Lemma 2.6 we get  $\phi_n^\alpha(a, b) = 0, \forall a, b \in U, \alpha \in \Gamma$  and  $n \in N$ .

**Lemma 2.11:** Let  $M$  be a 2-torsion free prime  $\Gamma$ -ring satisfying the condition (\*),  $U$  be an admissible Lie ideal of  $M$  and  $D = (d_i)_{i \in N}$  be a higher  $(U, M)$ -derivation of  $M$ . Then  $\phi_n^\alpha(u\beta u, m) = 0, \forall u \in U, m \in M, \alpha, \beta \in \Gamma$  and  $n \in N$ .

**Proof:** By Lemma 2.10, we have  $\phi_n^\alpha(u, v) = 0, \forall u, v \in U$  and  $\alpha \in \Gamma$ .

Replacing  $v$  by  $u\beta m - m\beta u$ , we get

$$\begin{aligned} 0 = \phi_n^\alpha(u, u\beta m - m\beta u) = d_n(u\alpha u\beta m) - d_n(u\alpha m\beta u) - \sum_{i+j=n} d_i(u)\alpha d_j(u\beta m - m\beta u) = \\ d_n(u\alpha u\beta m) - d_n(u\alpha m\beta u) - \sum_{i+j=n} (d_i(u)\alpha \sum_{p+q=j} d_p(u)\beta d_q(m) - d_p(m)\beta d_q(u)) = \\ d_n(u\alpha u\beta m) - d_n(u\alpha m\beta u) - \sum_{i+p+q=n} d_i(u)\alpha d_p(u)\beta d_q(m) + \\ \sum_{i+p+q=n} d_i(u)\alpha d_p(m)\beta d_q(u) = d_n(u\alpha u\beta m) - \sum_{s+q=n} (\sum_{i+p=s} d_i(u)\alpha d_p(u))\beta d_q(m) = \\ \phi_n^\alpha(u\beta u, m). \end{aligned}$$

### 3. $(U, M)$ -DERIVATION IN COMPLETELY SEMIPRIME $\Gamma$ -RING

Now we prove our main result in the following way:

**Theorem 3.1:** Let  $M$  be a 2-torsion free completely semiprime  $\Gamma$ -ring satisfying the condition (\*),  $U$  be an admissible Lie ideal of  $M$  and  $D = (d_i)_{i \in \mathbb{N}}$  be a higher  $(U, M)$ -derivation of  $M$ . Then  $d_n(u\alpha m) = \sum_{i+j=n} d_i(u)\alpha d_j(m)$ ,  $\forall u \in U, m \in M, \alpha \in \Gamma$  and  $n \in \mathbb{N}$ .

**Proof:** By the definition  $\phi_0^\alpha(u, m) = 0, \forall u \in U, m \in M, \alpha \in \Gamma$ .

Also by Theorem 2.1,  $\phi_1^\alpha(u, m) = 0, \forall u \in U, m \in M, \alpha \in \Gamma$ .

Now we proceed by induction as

Suppose that  $\phi_p^\alpha(u, m) = 0, \forall u \in U, m \in M, \alpha \in \Gamma$  and  $p \in \mathbb{N}$

This implies that

$$d_p(u\alpha m) = \sum_{i+j=p} d_i(u)\alpha d_j(m), u \in U, m \in M \text{ and } \alpha \in \Gamma \text{ and } p < n, \text{ where } p, n \in \mathbb{N}$$

Since  $D = (d_i)_{i \in \mathbb{N}}$  is a higher  $(U, M)$ -derivation of  $M$ , thus we have

$$\begin{aligned} d_n(u\alpha(u\beta m) + (u\beta m)\alpha u) &= \sum_{i+j=n} d_i(u)\alpha d_j(u\beta m) + d_i(u\beta m)\alpha d_j(u) = u\alpha d_n(u\beta m) + \\ d_n(u)\alpha(u\beta m) &+ \sum_{i+j=n}^{i,j < n} d_i(u)\alpha d_j(u\beta m) + (u\beta m)\alpha d_n(u) + d_n(u\beta m)\alpha(u) + \\ \sum_{i+j=n}^{i,j < n} d_i(u\beta m)\alpha d_j(u) &= u\alpha d_n(u\beta m) + d_n(u)\alpha(u\beta m) + \\ \sum_{i+j=n}^{i,j < n} d_i(u)\alpha \sum_{s+t=j} d_s(u)\beta d_t(m) &+ (u\beta m)\alpha d_n(u) + d_n(u\beta m)\alpha(u) + \\ \sum_{i+j=n}^{i,j < n} \sum_{l+q=i} d_l(u)\beta d_q(m)\alpha d_j(u) &= u\alpha d_n(u\beta m) + d_n(u)\alpha(u\beta m) + \\ \sum_{i+s+t=n}^{i,s+t < n} d_i(u)\alpha d_s(u)\beta d_t(m) &+ (u\beta m)\alpha d_n(u) + d_n(u\beta m)\alpha(u) + \\ \sum_{l+q+j=n}^{l+q,j < n} d_l(u)\beta d_q(m)\alpha d_j(u) & \end{aligned} \quad (7)$$

On the other hand, using Lemma 2.7 and 2.11, we get

$$\begin{aligned} d_n(u\alpha(u\beta m) + (u\beta m)\alpha u) &= d_n(u\alpha u\beta m) + d_n(u\beta m\alpha u) = \sum_{i+j=n} d_i(u\alpha u)\beta d_j(m) + \\ \sum_{i+j+k=n} d_i(u)\beta d_j(m)\alpha d_k(u) &= \sum_{p+q+j=n} d_p(u)\alpha d_q(u)\beta d_j(m) + \\ \sum_{i+j+k=n} d_i(u)\beta d_j(m)\alpha d_k(u) &= d_n(u)\alpha u\beta m + u\alpha \sum_{q+j=n} d_q(u)\beta d_j(m) + \\ \sum_{p+q+j=n}^{p,q+j < n} d_p(u)\alpha d_q(u)\beta d_j(m) &+ u\beta m\alpha d_n(u) + \sum_{i+j=n} d_i(u)\beta d_j(m)\alpha u + \\ \sum_{i+j+k=n}^{i+j,k < n} d_i(u)\beta d_j(u)\alpha d_k(m) & \end{aligned} \quad (8)$$

By comparing (7) and (8) and using the condition (\*), we get

$$\phi_n^\alpha(u, m)\beta u + u\beta \phi_n^\alpha(u, m) = 0, \forall u \in U, m \in M, \alpha, \beta \in \Gamma \text{ and } n \in \mathbb{N} \quad (9)$$

Linearizing of (9) with respect to  $u$ , gives us

$$\phi_n^\alpha(u, m)\beta v + \phi_n^\alpha(v, m)\beta u + u\beta\phi_n^\alpha(v, m) + v\beta\phi_n^\alpha(u, m) = 0, \forall u, v \in U, m \in M, \alpha, \beta \in \Gamma \text{ and } n \in N$$

Replacing  $v$  by  $v\alpha v$  and using Lemma 2.4, we get  $\phi_n^\alpha(u, m) = 0, \forall u \in U, m \in M, \alpha \in \Gamma$  and  $n \in N$ . Consequently, we get  $d_n(u\alpha m) = \sum_{i+j=n} d_i(u)\alpha d_j(m), \forall u \in U, m \in M, \alpha \in \Gamma$  and  $n \in N$ .

## REFERENCES

- [1] W.E. Barnes; "On the  $\Gamma$ -rings of Nobusawa", Pacific J. Math., Vol. 18(3), pp. 411-422, 1966.
- [2] N. Nobusawa; "On the generalization of the ring theory", Osaka J. Math., Vol. 1, pp. 81-89, 1964.
- [3] I.N. Herstein; "Jordan derivations of prime rings", Proc. Amer. Math. Soc., Vol. 8, pp. 1104-1110, 1957.
- [4] A.K. Faraj, C. Haetinger and A.H. Majeed; "Generalized Higher (U, R)-Derivations", JP Journal of Algebra, Vol. 16(2), pp. 119-142, 2010.
- [5] R. Awtar; "Lie ideals and Jordan derivations of prime rings", Proc. Amer. Math. Soc., Vol. 90(1), pp. 9-14, 1984.
- [6] M. Sapanci and A. Nakajima; "Jordan Derivations on Completely Prime  $\Gamma$ -Rings", Math. Japonica, Vol. 46(1), pp. 47-51, 1997.
- [7] K.W. Jun and B.D. Kim; "A note on Jordan left derivations", Bull. Korean Math. Soc., Vol. 33(2), pp. 221-228, 1996.
- [8] Y. Ceven; "Jordan left derivations on completely prime gamma rings", C.U. Fen-Edebiyat Fakultesi Fen Bilimleri Dergisi, Vol. 23, pp. 39-43, 2002.
- [9] A.K. Halder and A.C. Paul; "Jordan left derivations on Lie ideals of prime  $\Gamma$ -rings", Punjab University J. Math., Vol. 44, pp. 23-29, 2012.
- [10] M.M. Rahman and A.C. Paul; Generalized higher (U,M)-Derivations in Prime  $\Gamma$ -Rings", Bangladesh J. of Sci. Res., Vol. 27(2), pp. 143-153, 2014.